

**Topic : Quadratic Equation**

**Type of Questions**

**M.M., Min.**

Comprehension (no negative marking) Q.1 to 3

(3 marks, 3 min.)

[9, 9]

Single choice Objective (no negative marking) Q.4,5,6,7

(3 marks, 3 min.)

[12, 12]

Subjective Questions (no negative marking) Q.8


(4 marks, 5 min.)

[4, 5]

**COMPREHENSION (Q.No. 1 to 3)**

Consider the equation  $|x^2 - 2x - 3| = m, m \in \mathbb{R}$

- If the given equation has four solutions, then  
 (A)  $m \in (0, \infty)$       (B)  $m \in (-1, 3)$       (C)  $m \in (0, 4)$       (D) none of these
- If the given equation has three solutions, then  
 (A)  $m \in (0, \infty)$       (B)  $m \in \{4\}$       (C)  $m \in (0, 4)$       (D)  $m \in (-1, 3)$
- If the given equation has two solutions, then  
 (A)  $m \in [4, \infty)$       (B)  $m \in (-1, 3)$       (C)  $m \in (4, \infty) \cup \{0\}$       (D)  $m = 0$
- Let a, b, c be three roots of the equation  $x^3 + x^2 - 333x - 1002 = 0$ , then  $(\sum a^3) - 2 \sum a$  is equal to  
 (A) 2008      (B) 2000      (C) 2006      (D) 2002
- Number of real solutions of the equation  $x^2 + \left(\frac{x}{x-1}\right)^2 = 8$  is  
 (A) 3      (B) 4      (C) 6      (D) 0
- If  $y = ax^2 + bx + c$  represents the curve given in the figure and  $b^2 = 2(b + 2ac)$ , where  $a \neq 0$  and  $AP = 3$  units, then  $OP =$



- $\frac{3}{2}$       (B)  $\frac{3}{4}$       (C) 3      (D) 6
- If  $mx^2 - 9mx + 5m + 1 > 0, \forall x \in \mathbb{R}$ , then m lies in the interval  
 (A)  $\left(-\frac{4}{61}, 0\right)$       (B)  $\left[0, \frac{4}{61}\right)$       (C)  $\left(\frac{4}{61}, \frac{61}{4}\right)$       (D)  $\left(-\frac{61}{4}, 0\right)$
- Find the range of values of 'a' such that  $f(x) = \frac{ax^2 + 2(a+1)x + 9a + 4}{x^2 - 8x + 32}$  is always negative?

## Answers Key

1. (C)    2. (B)    3. (C)    4. (A) 5. (A)

6. (C)    7. (B)    8. a  $\left(-\infty, -\frac{1}{2}\right)$