

- **Inequalities:** Two real numbers or two algebraic expressions related by the symbols $<$, $>$, \leq or \geq form an inequality. For example: $3x < 20$, $4x + y < 12$.
- Equal numbers may be added to (or subtracted from) both sides of an inequality.
- Both sides of an inequality can be multiplied (or divided) by the same positive number. But when both sides are multiplied (or divided) by a negative number, then the inequality is reversed.
- **Linear Inequality:** An inequality is said to be linear, if each variable occurs in first degree only and there is no term involving the product of the variables. For example: $ax + b < 0$, $ax + b \leq 0$, $ax + by + c > 0$, $ax + by + c \geq 0$.
- **Closed Interval:** If a and b are real numbers, such that $a < b$, then the set of all real numbers x such that $a \leq x \leq b$ is called a closed interval and is denoted by $[a, b]$. Therefore, $[a, b] = \{x : a \leq x \leq b, x \in \mathbb{R}\}$.
- **Open Interval:** If a and b are real numbers, such that $a < b$, then the set of all real numbers x such that $a < x < b$ is called a open interval and is denoted by (a, b) or $]a, b[$. Therefore, $(a, b) = \{x : a < x < b, x \in \mathbb{R}\}$.
- **Solution of an Inequality:** The values of x , which make an inequality a true statement, are called solutions of the inequality.
- To represent $x < a$ (or $x > a$) on a number line, put a circle on the number and dark line to the left (or right) of the number a .
- To represent $x \leq a$ (or $x \geq a$) on a number line, put a dark circle on the number and dark the line to the left (or right) of the number x .
- If an inequality is having \leq or \geq symbol, then the points on the line are also included in the solutions of the inequality and the graph of the inequality lies left (below) or right (above) of the graph of the equality represented by dark line that satisfies an arbitrary point in that part.

- If an inequality is having $<$ or $>$ symbol, then the points on the line are not included in the solutions of the inequality and the graph of the inequality lies to the left (below) or right (above) of the graph of the corresponding equality represented by dotted line that satisfies an arbitrary point in that part.
- **To remove the denominator when the sign of the value of the denominator is unknown, may be +ve or -ve:** We multiply by the square of the denominator. Square of the denominator is always positive. On multiplication of an inequation by a positive number the sign of inequality does not change.
- **Solution of a system of Linear Inequality:** The solution set of a system of linear inequality in one variable is defined as the intersection of the solution sets of the linear inequality of the system. For example: If we have two solution sets $x > 4$ and $x > 6$, then solution of the system is the intersection of $x > 4$ and $x > 6$, i.e., $x > 6$.